

DUPIXENT[®] 
(dupilumab) Injection 300mg

HAVE DIFFICULTY SWALLOWING?

DU MORE TO HELP EASE
E OE
(EOSINOPHILIC ESOPHAGITIS)

A breakthrough treatment in EoE for adults and children 12+ years of age who weigh at least 88 lb (40 kg)

INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 12 years of age and older, who weigh at least 88 pounds (40 kg), with eosinophilic esophagitis (EoE). It is not known if DUPIXENT is safe and effective in children with eosinophilic esophagitis under 12 years of age and who weigh at least 88 pounds (40 kg).

IMPORTANT SAFETY INFORMATION


Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.


WHAT IS EoE?

Eosinophilic Esophagitis (EoE) is a chronic and progressive condition that can damage the esophagus over time.


People with EoE have inflammation in the esophagus, the tube that connects the throat to the stomach. The inflammation occurs when a large number of immune cells (including eosinophils) build up within the esophagus. This buildup is associated with an abnormal response of the immune system, known as type 2 inflammation.



Normal esophagus



Immune cell buildup causes inflammation



Inflammation leads to difficulty swallowing and other symptoms of EoE

For some people, inflammation and the associated changes in the esophagus could lead to food getting stuck in the esophagus, also known as food impaction. This can result in a trip to the ER.

EoE can be mistaken for other conditions, such as food allergies or GERD (gastroesophageal reflux disease). It's important for you or your loved one to recognize the physical symptoms and coping clues of EoE, such as difficulty swallowing and excessive chewing.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

WHO GETS EoE?

EoE can develop in children or adults. Symptoms vary by age:

- Children (ages 12-17) may have difficulty swallowing, food impaction, and abdominal pain, and vomit
- Adults may have difficulty swallowing and food impaction



HOW IS EoE DIAGNOSED?

A Gastroenterologist (GI) needs to conduct tests, including an "endoscopy with biopsy," to determine whether there's a high eosinophil count.

If you're currently seeing another specialist, such as an Allergist or a Pediatrician, or a PCP (Primary Care Practitioner), and suspect you may have EoE, you may need a referral to see a GI.

A BREAK-THROUGH TREATMENT FOR EoE

UNTIL DUPIXENT, THERE WAS NO FDA-APPROVED WAY TO TREAT EoE.

PPIs and topical swallowed corticosteroids are prescribed to help manage the disease and may provide temporary relief. DUPIXENT is not a PPI or a topical swallowed corticosteroid. It targets an underlying source of the inflammation that contributes to EoE.

PPIs and topical swallowed corticosteroids are not FDA-approved to treat EoE.



AT 24 WEEKS, PEOPLE TAKING DUPIXENT EXPERIENCED



INCREASED EASE OF SWALLOWING*

That means less choking, chest pain, or food getting stuck in the esophagus.



DECREASED INFLAMMATION IN THE ESOPHAGUS*

DUPIXENT reduced signs of inflammation.

*Based on clinical trials with 240 adult and pediatric patients (12+ years of age, weighing at least 88 lb) with EoE at week 24.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

THE FIRST AND ONLY TREATMENT OF ITS KIND

DUPIXENT is a biologic that helps block a key source of inflammation inside the body that can cause EoE.

Biologics are processed in the body differently than oral or topical medications. Most biologic medications come as liquids and are administered by injection.

DUPIXENT can be administered either by a doctor or at home after proper training.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have EoE and asthma and use an asthma medicine. **Do not** change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).



HAVE DIFFICULTY SWALLOWING?

**DU MORE
TO HELP EASE
EOE**

TALK TO YOUR DOCTOR

Bring this with you to your next doctor visit to help discuss if DUPIXENT is right for you or your loved one.

It's important to know the signs and symptoms of EoE. Check all that you are currently experiencing.

SYMPTOMS

- Difficulty swallowing (dysphagia)
- Food stuck in your throat (impaction)
- Coughing, choking, or vomiting
- Chest pain/heartburn, even after taking proton pump inhibitors (PPIs)

COPING MECHANISMS

- Chopping food into tiny pieces
- Mixing food with sauces or dips
- Eating slowly, chewing excessively
- Drinking a lot of liquids with food
- Avoiding certain foods
- Avoiding social occasions that center around eating

IMPORTANT SAFETY INFORMATION (CONT'D)

DUPIXENT can cause serious side effects, including:

- **Allergic reactions. DUPIXENT can cause allergic reactions that can sometimes be severe.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

HOW OFTEN DO YOU USE THESE COPING MECHANISMS?

- Daily Several times a week Once or twice a month

ARE YOU CURRENTLY TAKING ANY MEDICATIONS TO HELP MANAGE YOUR SYMPTOMS?

- No
- Yes. **If so**, what are you taking? _____

QUESTIONS TO ASK YOUR DOCTOR

How could DUPIXENT help me or my teen?

What results have you seen with EoE patients on DUPIXENT?

What support is available for patients taking DUPIXENT?

Use the space below to write your own questions or any notes.

HOW DUPIXENT[®] (dupilumab) WORKS

DUPIXENT targets an underlying source of type 2 inflammation and

- Is not a steroid
- Is not an immunosuppressant
- Is given subcutaneously (under the skin)

HOW DUPIXENT IS TAKEN

DUPIXENT is a weekly single-dose injection that can be given by your doctor in an office or a clinic, or can be taken at home. If your healthcare provider decides that you or a caregiver can give DUPIXENT injections, you or your caregiver should receive training on the right way to prepare and inject DUPIXENT. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult.



PRE-FILLED PEN

A hidden needle means you or your caregiver won't have to see the needle, and a viewing window and 2 click sounds let you know when your injection is complete.



PRE-FILLED SYRINGE

A contoured finger grip provides a firm hold and syringe design lets you or your caregiver control injection speed. The device features a needle shield.

QUESTIONS ABOUT INJECTING?



If you have questions about taking DUPIXENT, we're here for you. Scan or [CLICK HERE](#) for supplemental injection resources, helpful tips, and training videos.

IMPORTANT SAFETY INFORMATION (CONT'D)

DUPIXENT can cause serious side effects, including:

- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

The most common side effects in patients with eosinophilic esophagitis include injection site reactions, upper respiratory tract infections, cold sores in your mouth or on your lips, and joint pain (arthralgia).

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

HAS YOUR SPECIALIST PRESCRIBED DUPIXENT[®] (dupilumab)?

Your specialist is your best resource for information about DUPIXENT and should be your primary point of contact throughout your treatment journey. The following information describes how *DUPIXENT MyWay* can provide additional support resources and help you receive DUPIXENT as quickly as possible.

Your specialist prescribes DUPIXENT, and you enroll into *DUPIXENT MyWay*.

- You or your specialist can download the enrollment form on [DUPIXENT.com/support-savings/DUPIXENT-my-way](https://www.dupixent.com/support-savings/dupixent-my-way). Then fill it out and fax back to *DUPIXENT MyWay* at 1-844-387-9370
- You or your specialist can call 1-844-DUPIXEN(T), option 1
- Providing your email address allows *DUPIXENT MyWay* to give you more support resources about DUPIXENT

IMPORTANT SAFETY INFORMATION (CONT'D)

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

WHAT CAN *DUPIXENT MyWay*[®] DO FOR YOU?

DUPIXENT MyWay is a patient support program designed to help you stay on track throughout your treatment journey.

FOR ELIGIBLE PATIENTS, *DUPIXENT MyWay* CAN



Help research and explain your insurance benefits



Find ways to help with the out-of-pocket cost of DUPIXENT



Give you access to one-on-one support for DUPIXENT



Provide supplemental injection training in person, virtually, or over the phone



Help you schedule deliveries of your prescription

NEED HELP PAYING? \$0* COPAY MAY BE AVAILABLE

Enroll for the *DUPIXENT MyWay Copay Card*.

With the *DUPIXENT MyWay Copay Card*, eligible commercially insured patients may pay as little as \$0* copay per fill of DUPIXENT. And if you're eligible, you can download your card today. The program has an annual maximum of \$13,000.

You may be eligible for the copay card program if you

- Have commercial insurance, including health insurance exchanges, federal employee plans, or state employee plans
- Are a resident of the 50 United States, District of Columbia, or Puerto Rico
- Are prescribed DUPIXENT for an indication approved by the US Food and Drug Administration

DON'T HAVE INSURANCE?

There are other options for support. A *DUPIXENT MyWay Case Manager* can help qualified patients explore options to help cover the cost of DUPIXENT. If you need help paying for your prescription, the *DUPIXENT MyWay Patient Assistance Program* may be able to help.



Scan or click to go to [DUPIXENT.com/support-savings/DUPIXENT-my-way](https://www.dupilumab.com/support-savings/DUPIXENT-my-way) for full program information.

*Approval is not guaranteed. Program has an annual maximum of \$13,000. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed or restricted. *DUPIXENT MyWay* reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.

SUPPORT YOU DESERVE

The *DUPIXENT MyWay Copay Card* may help eligible patients cover the out-of-pocket cost of DUPIXENT.



ACTIVATING YOUR COPAY CARD

You can activate your Copay Card in 2 ways:

- Call 1-855-726-4702. Make sure you have the group number and ID numbers found on the Copay Card attached to this page
- Visit [DUPIXENT.com](https://www.dupilumab.com). You can apply and activate the card online

Once approved, provide the Copay Card numbers on your card to the specialty pharmacy when they call you to set up delivery of DUPIXENT. The pharmacy will apply the card to reduce your out-of-pocket costs and will note the card number in your record for future refills.

HAVE YOU ALREADY PAID OUT-OF-POCKET COSTS FOR DUPIXENT?

You may be eligible for reimbursement of your out-of-pocket expenses if you meet program eligibility requirements and

- You were enrolled in the Copay Card Program but did not use it when you filled your DUPIXENT prescription
- You paid a copay for DUPIXENT before enrolling into the Copay Card Program
- Your health plan did not accept your Copay Card

TO APPLY FOR REIMBURSEMENT

- Visit www.patientrebateonline.com to submit for reimbursement of out-of-pocket costs or to check the card balance. You will be asked to register and login
- Be sure to have ready
 - Your prescription insurance card
 - Prescription label
 - Your prescription receipt

Please allow 7-10 days to process your request.

Please see additional [Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.](#)

DUPIXENT[®]
(dupilumab) Injection 300mg

HAVE DIFFICULTY SWALLOWING?

DU MORE TO HELP EASE
EOE

ASK YOUR DOCTOR ABOUT DUPIXENT FOR EoE

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

Please see additional **Important Safety Information** throughout and accompanying full **Prescribing Information** including **Patient Information**.



To learn more, scan or click to go to
[DUPIXENT.com/eoe](https://www.dupilumab.com/eoe).

sanofi | **REGENERON**[®]

© 2023 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. DUPIXENT[®] and DUPIXENT MyWay[®]
are registered trademarks of Sanofi Biotechnology.
DUP.23.03.0199 05/23

Health information contained herein is provided for general educational purposes only. Your doctor is the single best source of information regarding your health. Please consult your doctor if you have any questions about your health or treatment.